



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2002

**Erratum: "Estimates on periodic and Dirichlet eigenvalues for the
Zakharov-Shabat system" [Asymptot. Anal. 25 (2001), no. 3-4, 201–237]**

Grébert, B ; Kappeler, T

Posted at the Zurich Open Repository and Archive, University of Zurich
ZORA URL: <https://doi.org/10.5167/uzh-21951>
Journal Article

Originally published at:

Grébert, B; Kappeler, T (2002). Erratum: "Estimates on periodic and Dirichlet eigenvalues for the Zakharov-Shabat system" [Asymptot. Anal. 25 (2001), no. 3-4, 201–237]. Asymptotic Analysis, 29(2):183.

Erratum

Estimates on periodic and Dirichlet eigenvalues for the Zakharov–Shabat system

by B. Grébert and T. Kappeler

[*Asymptotic Analysis* **25**(3,4) (2001), 201–237]

It was brought to our attention that inequality (A.6) on p. 232 in the proof of Proposition A.1 is wrong and therefore the proof for the bound (A.8) not correct.

The proof of Proposition A.1 can be easily corrected. Define for $n \geq 1$ the union of contours

$$\mathcal{R}_n = \left\{ \lambda \in \mathbb{C} \mid |\lambda| = n\pi - \frac{\pi}{2} \right\} \cup \left(\bigcup_{|k| \geq n} \left\{ \lambda \in \mathbb{C} \mid |\lambda - k\pi| = \frac{\pi}{2} \right\} \right).$$

For any $\lambda \in \bigcup_{n \geq 1} \mathcal{R}_n$, one has

$$L - \lambda = (\text{Id} + Q_\lambda)(L_0 - \lambda),$$

where Q_λ is the bounded operator on L^2 given by

$$Q_\lambda = \begin{pmatrix} 0 & \psi_1 \\ \psi_2 & 0 \end{pmatrix} (L_0 - \lambda)^{-1}.$$

To prove that $L - \lambda$ is invertible for any given λ in $\bigcup_{n \geq 1} \mathcal{R}_n$ it suffices to show that $\text{Id} + Q_\lambda$ is invertible. As in Section 2.2 we notice that whenever $\text{Id} - Q_\lambda^2$ is invertible one has

$$\text{Id} = (\text{Id} + Q_\lambda) \circ (\text{Id} - Q_\lambda) \circ (\text{Id} - Q_\lambda^2)^{-1}, \quad (\text{A.5}^*)$$

which implies that $\text{Id} + Q_\lambda$ is invertible. Using the assumption that w is a δ -weight one concludes from Lemmas 2.1 and 2.2 that there exists $N \geq 1$ so that for any $\lambda \in \bigcup_{n \geq N} \mathcal{R}_n$ and any 1-periodic potentials $\psi_1, \psi_2 \in H^w$ with $\|\psi_j\|_w \leq M$ one has

$$\|Q_\lambda^2\|_{\mathcal{L}(L^2)} \leq \frac{1}{2}. \quad (\text{A.8}^*)$$

The proof of Proposition A.1 can then be finished as in Appendix A and Lemma A.2 follows, using (A.5*) and (A.8*).